

The matrix square root

Next (and last, for us) matrix function: $A^{1/2}$, principal square root.

$A^{1/2}$ is well defined unless A has:

- ▶ Real eigenvalues $\lambda_i < 0$, or
- ▶ Non-trivial Jordan blocks at $\lambda_i = 0$ (because $g(x) = x^{1/2}$ is not differentiable).

Condition number / sensitivity

The Fréchet derivative of $f(X) = X^2$ is

$$L_{f,X}(E) = XE + EX, \quad \hat{L} = I \otimes X + X^T \otimes I.$$

The Fréchet derivative of $g(Y) = Y^{1/2}$ is its inverse,

$$\hat{L}_{g,Y} = (I \otimes Y^{1/2} + (Y^{1/2})^T \otimes I)^{-1}$$

with eigenvalues $\frac{1}{\lambda_i^{1/2} + \lambda_j^{1/2}}$, $i, j = 1, \dots, n$.

In particular, g is ill-conditioned for matrices that either:

- ▶ have a small eigenvalue (taking $i = j$), or
- ▶ have two complex conjugate eigenvalues close to the negative real axis (because then $\lambda_i^{1/2} \approx ai$, $\lambda_j^{1/2} \approx -ai$).

Schur method

Recall: Schur method:

1. Reduce to a triangular T using a Schur form;
2. Compute diagonal of $S = f(T)$;
3. Compute off-diagonal entries from $ST = TS$
Involves a denominator $t_{ii} - t_{jj}$: if it is 0, we must work on blocks.

In the case of $A^{1/2}$, we can use $S^2 = T$ to get the off-diagonal entries instead:

$$s_{ii}s_{jj} + s_{i,i+1}s_{i+1,j} + \cdots + s_{ij}s_{jj} = t_{ij}.$$

Involves a denominator $s_{ii} + s_{jj}$: always invertible because $s_{ii} + s_{jj} \in RHP$.

(This is what Matlab uses, by the way.)

Newton method

Newton method on $X^2 - A$:

$$X_{k+1} = X_k - E, \quad \text{where } E \text{ solves } EX_k + X_kE = X_k^2 - A.$$

Much more expensive than the Schur method: we solve a Sylvester equation at each step (and this requires a Schur form).

Trick: If X_0 commutes with A (for instance, taking $X_0 = \alpha I$), then $E = (2X_0)^{-1}(X_0^2 - A)$ and E, X_1 commute with A , too, ...

Resulting iteration:

(Modified) Newton iteration

$$X_{k+1} = \frac{1}{2}(X_k + X_k^{-1}A), \quad X_0 = \alpha I.$$

At each step, $X_k A = A X_k$.

Square root and sign

(Modified) Newton iteration

$$X_{k+1} = \frac{1}{2}(X_k + X_k^{-1}A), \quad X_0 = \alpha I.$$

Pre-multiply by $A^{-1/2}$, and use commutativity:

$$A^{-1/2}X_{k+1} = \frac{1}{2} \left(A^{-1/2}X_k + (A^{-1/2}X_k)^{-1} \right), \quad A^{-1/2}X_0 = \alpha A^{-1/2}.$$

This is the sign iteration! $A^{-1/2}X_k \rightarrow \text{sign}(A^{-1/2}) = I$.

Hence,

$X_k \rightarrow A^{1/2}$, i.e., the modified Newton iteration converges (for each starting point $X_0 = \alpha I$ with $\alpha > 0$).

Local convergence

True Newton

$$X_{k+1} = X_k - E, \quad \text{where } E \text{ solves } EX_k + X_k E = X_k^2 - A.$$

This is a Newton method, so it converges quadratically (locally).

Modified Newton

$$X_{k+1} = \frac{1}{2}(X_k + X_k^{-1}A).$$

The two iterations coincide, if $X_0 A = A X_0$in exact arithmetic!
In practice, this property is lost numerically. We need to study the convergence of MN separately.

MN is the fixed-point iteration associated to
 $h(X) = \frac{1}{2}(X + X^{-1}A).$

Local convergence

Local convergence of a fixed-point iteration depends on the eigenvalues of the Jacobian in the fixed-point.

The Jacobian / Fréchet derivative of $h(X) = \frac{1}{2}(X + X^{-1}A)$ is

$$L_{h,X}(E) = \frac{1}{2}(E + X^{-1}EX^{-1}A),$$

using $(X + E)^{-1} - X^{-1} = (X + E)^{-1}EX^{-1} = X^{-1}EX^{-1} + o(\|E\|)$.

Hence $L_{h,A^{1/2}} = \frac{1}{2}(E + A^{-1/2}EA^{1/2})$, or

$$\widehat{L}_{h,A^{1/2}} = \frac{1}{2} \left(I + (A^{1/2})^T \otimes A^{-1/2} \right).$$

It has eigenvalues $\frac{1}{2} + \frac{1}{2}\lambda_i^{1/2}\lambda_j^{-1/2}$, where λ_i are the eigenvalues of A .

It's easy to construct cases in which $L_{h,A^{1/2}}$ has eigenvalues with modulus > 1 , hence $A^{1/2}$ is an **unstable fixed point** of $h(X)$.